


QUEENSWAY

FACIAL AESTHETICS



smooth

facial anti-ageing treatments

Botox® Pre-treatment &
After Care Information

smooth



Botox® Treatment

Before Treatment:

- Treatment is not safe for people with certain medical conditions – be sure that you have informed your surgeon about your medical history including any medication you are taking.
- You should not go ahead with treatment if you are pregnant or breastfeeding.
- Please ensure you have fully discussed the proposed treatment with your surgeon, signed your consent form and you are happy to proceed.
- If you have any further questions, please ask before the treatment is carried out.

After Treatment:

- Slight redness, swelling, bruising and some tenderness after the treatment is common and should settle down within a few hours. A cold pack may help this.
- Occasionally headaches and eyelid drooping may occur – this is unusual and should resolve within a few weeks.
- Avoid facial beauty treatments and massage for the next 48 hours as you may cause unwanted spreading of the Botox into adjacent muscles.
- Avoid rubbing or pressing the area for the rest of the day – take care when applying make-up.
- The effects can take up to 10 days to fully work but should be apparent after a few days.
- You will usually be asked to come back for a review appointment 2 – 3 weeks after treatment to ensure that you are happy with the results.

Please contact the clinic if you have any concerns at all.

Queensway Dental Clinic, 293 Yarm Road, Darlington, DL1 1BA
Queensway Dental Clinic, 170 Queensway, Billingham, TS23 2NT

Tel: 01325 381 928
Tel: 01642 554 667

| www.queensway.co.uk